arXiv:1209.5188v2 [hep-ph] 25 Sep 2012

Zero angle Light-by light scattering tensor

A. I. Ahmadov[1,2] [*], S. Bakmaev[3] [†] and E. A. Kuraev[1] [‡]
[1] *Bogoliubov Laboratory of Theoretical Physics, Joint Institute for Nuclear Research, Dubna, 141980 Russia*
[2] *Institute of Physics, Azerbaijan National Academy of Sciences, Baku, Azerbaijan*
[3] *Dzhelepov Laboratory of Nuclear Problems, Joint Institute for Nuclear Research, Dubna, 141980 Russia*
(Dated: September 26, 2012)

We calculate the bilinear combination of Dirac tensors, describing the creation of a pair of charged particles (pi-meson or leptons) by virtual photons integrated on the final particles phase volume. It can be interpreted as a s channel discontinuity of the zero angle light-by light scattering tensor.

I. INTRODUCTION

A lot of attention was paid to the problem of describing the lowest order nonlinear effect of Quantum Electrodynamics-the scattering of photon on photon. One of earlier papers [1] reveals rather complicate structure of tensor describing the light by light scattering (lbl) process

$$\gamma^*(k_1,\mu)+\gamma^*(k_2,\nu)\to\gamma^*(k_3,\rho)+\gamma^*(k_4,\sigma) \tag{1}$$

with all photons off mass shell.

$$\begin{aligned}G_{\mu\nu\rho\sigma}(1234)=\sum_{24perm}\{&\frac{1}{8}A^{2143}(1234)k_\mu^{(2)}k_\nu^{(1)}k_\rho^{(4)}k_\sigma^{(3)}+\frac{1}{4}A^{2341}(1234)k_\mu^{(2)}k_\nu^{(3)}k_\rho^{(4)}k_\sigma^{(1)}+\\&\frac{1}{2}A^{2111}(1234)k_\mu^{(2)}k_\nu^{(1)}k_\rho^{(1)}k_\sigma^{(1)}+\frac{1}{2}A^{2121}(1234)k_\mu^{(2)}k_\nu^{(1)}k_\rho^{(2)}k_\sigma^{(1)}+\\&A^{2311}(1234)k_\mu^{(2)}k_\nu^{(3)}k_\rho^{(1)}k_\sigma^{(1)}+A^{2123}(1234)k_\mu^{(2)}k_\nu^{(1)}k_\rho^{(2)}k_\sigma^{(3)}+\\&\frac{1}{2}B_1^{11}(1234)\delta_{\mu,\nu}k_\rho^{(1)}k_\sigma^{(1)}+\frac{1}{2}B_1^{12}(1234)\delta_{\mu,\nu}k_\rho^{(1)}k_\sigma^{(2)}+\\&B_1^{13}(1234)\delta_{\mu,nu}k_\rho^{(1)}k_\sigma^{(3)}+\frac{1}{4}B_1^{43}(1234)\delta_{\mu,\nu}k_\rho^{(4)}k_\sigma^{(3)}+\frac{1}{8}C_1(1234)\delta_{\mu\nu}\delta_{\rho,\sigma}\},\end{aligned} \tag{2}$$

with notations given in [1]. A detailed investigation was done in papers of 1960-1980 years(see for instance [2] and references therein). Different channels with some of photons on mass shell was investigated using different approaches. One of the them was the photon splitting in Coulomb field of a heavy nuclei [3] was described using the double dispersion approach.

Nevertheless for many applications-the forward kinematics for instance the lbl tensor can be written in compact form see (15) below. It is the aim of our paper-to calculate the s channel discontinuity of lbl tensor with fermions or scalar particles intermediate state, which is associated with the scattering process

$$\gamma(k,\mu)+\gamma(q,\nu)\to\gamma(k,\mu_1)+\gamma(q,\nu_1), \tag{3}$$

namely

$$\Delta_s L^{\mu\nu;\mu_1\nu_1}(k,q;k,q)=(4\pi\alpha)^2\int T^{\mu\nu}(k,q;q_+,q_-)(T^{\mu_1\nu_1}(k,q;q_+,q_-))^*d\Gamma_2, \tag{4}$$

with the phase volume of the intermediate state with two on mass shell charged particles

$$\begin{aligned}d\Gamma_2=\frac{1}{(2\pi)^2}d^4q_-d^4q_+\delta(q_-^2-m^2)\delta(q_+^2-m^2)\delta^4(k+q-q_+-q_-)=\\\frac{1}{(2\pi)^2}\frac{d^3q_+}{2E_+}\frac{d^3q_-}{2E_-}\delta^4(k+q-q_+-q_-).\end{aligned} \tag{5}$$

---

[*] E-mail: ahmadov@theor.jinr.ru
[†] E-mail: bakmaev@jinr.ru
[‡] E-mail: kuraev@theor.jinr.ru

Writing down the phase volume in form (we will work in the center of mass reference frame $\vec{k}+\vec{q}=0$):

$$\int d\Gamma_2 = \frac{\beta}{8\pi}\frac{1}{2\pi}\int\limits_0^{2\pi} d\phi \frac{1}{2}\int\limits_{-1}^{1} dc, \tag{6}$$

with 4-vector q_- component defined as $q_- = (\sqrt{s}/2)[1, \beta\vec{n}]$, $\vec{n} = (c, \sin\theta\cos\varphi, \sin\theta\sin\varphi)$ and

$$s = (k+q)^2 = (q_+ + q_-)^2; \qquad \beta = \sqrt{1-\frac{4m^2}{s}}, c = \cos\theta. \tag{7}$$

Here $\theta = (\vec{k}, \vec{q_-})$ and ϕ-are the polar and azimuthal angles which define the direction of 3-vector $\vec{q_-}$ in frames with z axes directed along $\vec{k}$. The integration on phase volume is essentially the angular averaging:

$$\begin{aligned} \Delta_s L^{\mu\nu;\mu_1\nu_1}(k,q;k,q) &= (4\pi\alpha)^2\frac{\beta}{8\pi} l^{\mu\nu\mu_1\nu_1}, \\ l^{\mu\nu\mu_1\nu_1} &=< T^{\mu\nu}(k,q;q_+,q_-)(T^{\mu_1\nu_1}(k,q;q_+,q_-))^* >, \\ < F > &= \frac{1}{2\pi}\int\limits_0^{2\pi} d\phi\frac{1}{2}\int\limits_{-1}^{1} dc F(c,\phi). \end{aligned} \tag{8}$$

II. LIGHT-LIKE BASIS. GENERAL FORM OF THE ZERO-ANGLE SCATTERING TENSOR

It's convenient to introduce two linear combinations of 4-momenta of photons $k = (k_0, |\vec{k}|, 0, 0); q = (q_0, -|\vec{k}|, 0, 0)$

$$\chi = \frac{k+q}{\sqrt{s}} = (1,0,0,0), \qquad r = \frac{q_0 k - k_0 q}{|\vec{k}|\sqrt{s}} = (0,1,0,0)$$
$$\chi^2 = 1, \qquad r^2 = -1, \qquad (\chi r) = 0 \tag{9}$$

and the "transversal" metric tensor

$$g^{\perp}_{\mu\nu} = \delta_{\mu,3}\delta_{\nu,3} + \delta_{\mu,4}\delta_{\nu,4}. \tag{10}$$

It have only transversal to 4-vectors k, q component

$$g^{\perp}_{\mu\nu}k^{\mu} = g^{\perp}_{\mu\nu}q^{\mu} = 0. \tag{11}$$

Transversal tensor can be written as

$$g^{\perp}_{\mu\nu} = -g_{\mu\nu} + \chi_\mu\chi_\nu - r_\mu r_\nu. \tag{12}$$

The photon's momenta can be written in terms of 4-vectors χ, r as

$$k = k_0\chi + pr, \qquad q = q_0\chi - pr, \qquad p = \frac{1}{2}\sqrt{\frac{\Lambda}{s}}. \tag{13}$$

Explicit form of $k_0, q_0, |\vec{k}| = |\vec{q}|, \Lambda = \Lambda(s, k^2, q^2)$ are given in Appendix A.

Besides we introduce two vectors orthogonal to k, q:

$$\bar{k} = p\chi + k_0 r; \qquad \bar{q} = p\chi - q_0 r; \qquad (k\bar{k}) = 0; \qquad (q\bar{q}) = 0. \tag{14}$$

The $l^{\mu\nu\mu_1\nu_1}$ tensor can be written in terms of $g^{\perp}$ and $\bar{k}, \bar{q}$ which are explicit gauge-invariant quantities:

$$l^i_{\mu\nu\mu_1\nu_1} = a^i_F F + a^i_G G + a^i_H H + a^i_a a + a^i_b b + a^i_c c + a^i_d d + a^i_e e + a^i_f f + a^i_h \bar{k}_\mu\bar{k}_{\mu_1}\bar{q}_\nu\bar{q}_{\nu_1}, \qquad i = \pi, \mu, \tag{15}$$

with c-number coefficients a^i_j given below and tensor structures

$$\begin{aligned} F &= g^{\perp}_{\mu\mu_1}g^{\perp}_{\nu\nu_1}; & G &= g^{\perp}_{\mu\nu_1}g^{\perp}_{\nu\mu_1}; & H &= g^{\perp}_{\mu\nu}g^{\perp}_{\mu_1\nu_1}; \\ a &= g^{\perp}_{\mu\mu_1}\bar{q}_\nu\bar{q}_{\nu_1}; & b &= g^{\perp}_{\mu\nu}\bar{q}_{\nu_1}\bar{k}_{\mu_1}; & c &= g^{\perp}_{\nu\mu_1}\bar{q}_{\nu_1}\bar{k}_\mu; \\ d &= g^{\perp}_{\mu\nu_1}\bar{q}_\nu\bar{k}_{\mu_1}; & e &= g^{\perp}_{\nu_1\mu_1}\bar{q}_\nu\bar{k}_\mu; & f &= g^{\perp}_{\nu\nu_1}\bar{k}_\mu\bar{k}_{\mu_1}. \end{aligned} \tag{16}$$

III. RESULTS

For the case of pion pair production we obtain (details in Appendices A,B,C):

$$\begin{aligned}
a_H^\pi &=< 4+16\lambda\frac{1}{d}\beta_1+16\frac{1}{d^2}\lambda^2\beta_2 >;\\
a_G^\pi = a_F^\pi &= 16 < \frac{1}{d^2}\lambda^2\beta_2 >;\\
p^2a_b^\pi = p^2a_e^\pi &=< 4[\frac{\lambda}{d}-4\frac{\lambda^2}{d^2}]\beta_1-4+2\frac{\lambda}{d}[-5+\rho] >;\\
p^2a_d^\pi = p^2a_c^\pi &=< 4\lambda^2(\frac{1}{d}-\frac{1}{d^2})(\rho-1)\beta_1 >;\\
p^2a_a^\pi &=< 4\lambda^2\eta^2(\frac{1}{d}-\frac{1}{d^2})\beta_1 >;\\
p^2a_a^\pi &=< 4\lambda^2\sigma^2(\frac{1}{d}-\frac{1}{d^2})\beta_1 >;\\
p^4a_h^\pi &=< 4+4(1-\rho)\frac{\lambda}{d}+4[16+(1-\rho)^2]\frac{\lambda^2}{d^2} > .
\end{aligned} \tag{17}$$

The coefficients for tensor for the case of lepton pair in the intermediate state is

$$\begin{aligned}
p^2a_a^\mu &= \frac{8\lambda^2}{d^2}[(1-d)\epsilon^2+[\frac{1}{4}d(\delta-\epsilon-1)^2-\lambda^2(\delta+\epsilon-1)^2\epsilon]\beta_1];\\
p^2a_b^\mu &= -\frac{4(1-d)}{d^2}[\frac{1}{2}d-2\lambda^2\epsilon\delta+\lambda^2(\delta+\epsilon-1)^2\beta_1];\\
p^2a_c^\mu &= -\frac{4\lambda^2}{d^2}[2(1-d)\epsilon\delta+[\frac{1}{2}(2-d)(\delta-\epsilon-1)^2-(1-d)(\delta+\epsilon-1)-2\epsilon(2-d)]\beta_1];\\
a_F^\mu &= \frac{8\lambda^2(\epsilon+\delta-1)^2}{d^2}[\frac{1}{4}d-\epsilon\delta\lambda^2+(\delta+\epsilon)\lambda^2\beta_1-\lambda^2\beta_2];\\
a_G^\mu &= -\frac{8\lambda^2(\epsilon+\delta-1)^2}{d^2}[\frac{1}{4}d-\epsilon\delta\lambda^2+(\delta+\epsilon-1)\lambda^2\beta_1+\lambda^2\beta_2];\\
a_H^\mu &= \frac{8(1-d)}{d^2}[[\frac{1}{4}d-\epsilon\delta\lambda^2+(\delta+\epsilon-1)\lambda^2\beta_1](1-d)-\lambda^4(\epsilon+\delta-1)^2\beta_2];\\
p^4a_h^\mu &= \frac{8(1-d)}{d^2}[\frac{1}{4}d-\epsilon\delta\lambda^2],
\end{aligned} \tag{18}$$

besides

$$a_f^\mu = a_a^\mu(\epsilon\leftrightarrow\delta); p^2a_e^\mu = a_b^\mu; a_d^\mu = a_c^\mu. \tag{19}$$

Here we use the notations

$$\begin{aligned}
\epsilon &= \frac{k^2}{s}; \delta = \frac{q^2}{s}; \sigma = \frac{q_0}{\sqrt{s}}; \eta = \frac{k_0}{\sqrt{s}};\\
\lambda &= \frac{1}{\epsilon+\delta-1}; \rho = \frac{\Lambda}{s^2}, \beta_1 = \frac{1}{2}\beta^2(1-c^2); \beta_2 = \frac{1}{8}\beta^4(1-c^2)^2.
\end{aligned} \tag{20}$$

The relevant integrals needed within operation of averaging over polar angle are given in Appendix.

IV. CONCLUSION. THE CASE OF REAL PHOTONS

As well as the results are given in relativistic-invariant form the form of *lbl* tensor in any reference frame can be obtained.

For the case of both photons on mass shell the 4-vectors $\bar{k}, \bar{q}$ become the light-like ones. Using

$$p = \frac{\sqrt{s}}{2}; \rho = 1, \lambda = -1; \eta = \sigma = \frac{1}{2}, \epsilon = \delta = 0, \tag{21}$$

we obtain

$$\begin{aligned}
a^\pi_H &=< 4-16\frac{1}{d}\beta_1+16\frac{1}{d^2}\beta_2 >;\\
a^\pi_G = a^\pi_F &= 16 < \frac{1}{d^2}\beta_2 >;\\
\frac{s}{4}a^\pi_b = \frac{s}{4}a^\pi_e &=< -4\beta_1[\frac{1}{d}+4\frac{1}{d^2}]-4+8\frac{1}{d} >;\\
\frac{s}{4}a^\pi_d &= p^2a^\pi_c = 0;\\
\frac{s}{4}a^\pi_a = \frac{s}{4}a^\pi_f &=< \beta_1(\frac{1}{d}-\frac{1}{d^2}) >;\\
\frac{s^2}{16}a^\pi_h &= 4 < 1+16\frac{1}{d^2} > .
\end{aligned} \tag{22}$$

and

$$\begin{aligned}
a^\mu_F &=< \frac{8}{d^2}[\frac{1}{4}d-\beta_2 >;\\
a^\mu_G &=< -\frac{8}{d^2}[\frac{1}{4}d-\beta_1+\beta_2 >;\\
a^\mu_H &=< \frac{8}{d^2}[(1-d)(\frac{1}{4}d-\beta_1)-\beta_2 >;\\
a^\mu_d = a^\mu_a = -a^\mu_c &= -a^\mu_d = \frac{2}{d}\beta_1;\\
a^\mu_b = a^\mu_e &= \frac{4}{d^2}(1-d)(\frac{1}{2}d-\beta_1),\\
d &= 1-\beta^2c^2.
\end{aligned} \tag{23}$$

As an important test is the correspondence with cross sections of pair production in real photons collisions. Really

$$\frac{d\sigma^{\gamma\gamma\to a\bar{a}}}{dc} = \frac{\alpha^2}{2(s-4m^2)}g^{\mu\mu_1}g^{\nu\nu_1}l_{\mu\nu\mu_1\nu_1}. \tag{24}$$

We obtain for the case of a pair of charged scalar particles [4]

$$\frac{d\sigma}{dc} = \frac{\alpha^2}{2s}\beta[1-\frac{8m^2}{s}\frac{1}{d}+\frac{32m^2}{s^2}\frac{1}{d^2}], \tag{25}$$

and for fermion pair

$$\frac{d\sigma}{dc} = \frac{2\alpha^2\beta}{s}[\frac{1+\beta^2c^2}{d}+\frac{8m^2}{sd}-\frac{32m^4}{s^2d^2}]. \tag{26}$$

Constructing the combination $g^{\mu\mu_1}g^{\nu\nu_1}l_{\mu\nu\mu_1\nu_1} = 4a_F + 2(a_G + a_H)$ one can be convinced that these tests are fulfilled.

V. ACKNOWLEDGEMENTS

Two of us (E.K.,Yu.B.) are grateful to RFBR grants 10-02-01295a; 11-02-00112 for support.

Appendix A: The charged pi-meson pair intermediate state. Kinematics

Matrix element of subprocess have a form

$$M^{\gamma^*\gamma^*\to q^-q^+}_{scal} = 4\pi\alpha T^{\mu\nu}, \tag{A1}$$

with

$$\begin{aligned}
T^{\mu\nu} &= \frac{(2q_- - k)^\mu(-2q_+ + q)^\nu}{d_-} + \frac{(k-2q_+)_\mu(2q_- - q)_\nu}{d_+} - 2g_{\mu\nu},\\
d_- &= (q_- - k)^2 - m^2 = k^2 - 2q_-k;\;\; d_+ = (-q_+ + k)^2 - m^2 = k^2 - 2q_-q.
\end{aligned} \tag{A2}$$

One can be convinced in fulfilment of gauge-invariance condition $T^{\mu\nu}k_\mu = T^{\mu\nu}q_\nu = 0$.

It seems to be convenient to rewrite the Dirac tensor $T^{\mu\nu}$ as

$$
\begin{aligned}
T^{\mu\nu} &= Aq_-^\mu q_-^\nu - 2q_-^\mu C^\nu - 2q_-^\nu D^\mu + r_{\mu\nu} + 2g^\perp_{\mu\nu}; \\
A = \frac{1}{d_-} + \frac{1}{d_+};\quad C^\nu &= \left(\frac{2k+q}{d_-} + \frac{q}{d_+}\right)^\nu;\ \ D^\mu = \left(\frac{k}{d_-} + \frac{2q+k}{d_+}\right)^\mu; \\
r_{\mu\nu} &= \frac{1}{d_-}k^\mu(2k+q)^\nu + \frac{1}{d_+}q_\nu(2q+k)_\mu - 2\chi_\mu\chi_\nu + 2r_\mu r_\nu; \\
g^\perp_{\mu\nu} &= -g_{\mu\nu} + \chi_\mu\chi_\nu - r_\mu r_\nu,
\end{aligned}
\tag{A3}
$$

Using the energy-momentum conservation law

$$k + q = q_+ + q_-$$

we find

$$
\begin{aligned}
k_0 &= \frac{s + k^2 - q^2}{2\sqrt{s}},\ \ q_0 = \frac{s + q^2 - k^2}{2\sqrt{s}}; \\
\vec{k}^2 &= \frac{\Lambda}{4s}, \Lambda = s^2 + (k^2)^2 + (q^2)^2 - 2sk^2 - 2sq^2 - 2k^2q^2.
\end{aligned}
\tag{A4}
$$

Below we will express the lbl tensor in terms of $g^\perp$ and $\bar{k}, \bar{q}$ which are explicit gauge-invariant quantities.

So we have (see (9)):

$$\Delta_s L^{\mu\nu;\mu_1\nu_1}(k,q;k,q) = \frac{\beta}{8\pi}(4\pi\alpha)^2 < T^{\mu\nu}(T^{\mu_1\nu_1})^* > . \tag{A5}$$

Appendix B: The charged lepton pair intermediate state

Matrix element of subprocess have a form

$$M^{\gamma^*\gamma^*\to q^- q^+}_{q_+q_-} = 4\pi\alpha T^{\mu\nu}_{q_+q_-}, \tag{B1}$$

with

$$T^{\mu\nu}_{q_+q_-} = \bar{u}(q_-)[\gamma_\mu \frac{\hat{q}_- - \hat{k} + m}{d_-}\gamma_\nu + \gamma_\nu \frac{-\hat{q}_+ + \hat{k} + m}{d_-}\gamma_\mu]v(q_+).$$

Using the on mass shell conditions of leptons it can be written as

$$
\begin{aligned}
T^{\mu\nu}_{q_+q_-} &= \bar{u}(q_-)[Q_\mu\gamma_\nu - \frac{1}{d_-}\gamma_\mu\hat{k}\gamma_\nu + \frac{1}{d_+}\gamma_\nu\hat{k}\gamma_\mu]v(q_+), \\
Q_\mu &= 2(\frac{q_-}{d_-} - \frac{q_+}{d_+})_\mu.
\end{aligned}
\tag{B2}
$$

Rather tedious calculation leads to result given above.

Appendix C: details of calculations

The expression for denominators are

$$d_- = \frac{k^2 + q^2 - s}{2}(1 + bc),\ \ d_+ = \frac{k^2 + q^2 - s}{2}(1 - bc),\ \ b = \beta\frac{\sqrt{\Lambda}}{k^2 + q^2 - s},\ c = \cos\theta,\ \theta = \widehat{\vec{q}_-\vec{k}}. \tag{C1}$$

Product of two Dirac tensors for pion pair have a form (notations in (A3)):

$$
\begin{aligned}
T_{\mu\nu}T_{\mu_1\nu_1} = 16A^2(q_-q_-q_-q_-)^{\mu\mu_1\nu\nu_1} - 8AC^{\nu_1}(q_-q_-q_-)^{\mu\nu\mu_1} - 8AC^\nu(q_-q_-q_-)^{\mu\mu_1\nu_1} - 8AD^{\mu_1}(q_-q_-q_-)^{\mu\nu\nu_1} - \\
8AD^\mu(q_-q_-q_-)^{\nu\mu_1\nu_1} + 4Ar_{\mu_1\nu_1}(q_-q_-)^{\mu\nu} + 4Ar_{\mu\nu}(q_-q_-)^{\mu_1\nu_1} + \\
4C^\nu C^{\nu_1}(q_-q_-)^{\mu\mu_1} + 4D^\mu D^{\mu_1}(q_-q_-)^{\nu\nu_1} + 4C^\nu D^{\mu_1}(q_-q_-)^{\mu\nu_1} + 4C^{\nu_1}D^\mu(q_-q_-)^{\nu\mu_1} - \\
2C^\nu r_{\mu_1\nu_1}q_-^\mu - 2C^{\nu_1}r_{\mu\nu}(q_-)^{\mu_1} - 2D^\mu r_{\mu_1\nu_1}q_-^\nu - 2D^{\mu_1}r_{\mu\nu}q_-^{\nu_1} + r_{\mu\nu}r_{\mu_1\nu_1} + 4g_\perp^{\mu\nu}q_\perp^{\mu_1\nu_1} + \\
2g_\perp^{\mu\nu}[4A(q_-q_-)^{\mu_1\nu_1} - 2C^{\nu_1}q_-^{\mu_1} - 2D^{\mu_1}q_-^{\nu_1} + r_{\mu_1\nu_1}] + 2g_\perp^{\mu_1\nu_1}[4A(q_-q_-)^{\mu\nu} - 2q_-^\mu C^\nu - 2q_-^\nu D^\mu + r_{\mu\nu}].
\end{aligned}
\tag{C2}
$$

The averaging of the relevant 4-vector product give

$$
\begin{aligned}
< q_-^\mu > &= \frac{\sqrt{s}}{2} < (\chi + \beta c r)^\mu >; \\
< q_-^\mu q_-^\nu > &= \frac{s}{4} < [\chi^2 + \beta c(\chi r) + \beta^2(c^2 r^2 + \frac{1}{2}(1-c^2) g_\perp)]^{\mu\nu} >; \\
< q_-^\mu q_-^\nu q_-^\lambda > &= \left(\frac{\sqrt{s}}{2}\right)^3 < \{\chi^3 + (\chi^2 r)\beta c + \beta^2[(\chi r^2) c^2 + \frac{1}{2}(1-c^2)(\chi g_\perp)] + \beta^3[c^3 r^3 + \frac{1}{2} c(1-c^2)(r g_\perp)]\}^{\mu\nu\lambda} >; \\
< q_-^\mu q_-^\nu q_-^\lambda q_-^\sigma > &= \left(\frac{s}{4}\right)^2 < \{\chi^4 + \beta c(\chi^3 r) + \\
&\quad \beta^2[c^2(\chi^2 r^2) + \frac{1}{2}(1-c^2)(\chi^2 g_\perp)] + \beta^3[c^3(\chi r^3) + \frac{1}{2} c(1-c^2)(\chi r g_\perp)] + \\
&\quad \beta^4[c^4 r^4 + \frac{1}{2}(c^2(1-c^2))(r^2 g_\perp) + \frac{1}{8}(1-c^2)^2 (g_\perp g_\perp)]\}^{\mu\nu\mu_1\nu_1} > . \qquad \text{(C3)}
\end{aligned}
$$

Here we imply

$$
\begin{aligned}
(ab)^{\mu\nu} = a^\mu b^\nu + a^\nu b^\mu; (a^n)^{\mu_1 \cdots \mu_n} = a^{\mu_1} ... a^{\mu_n}; (a^2 b)^{\mu_1\mu_2\mu_3} = a^{\mu_1} a^{\mu_2} b^{\mu_3} + ... + b^{\mu_1} a^{\mu_2} a^{\mu_3}; \\
(a g_\perp)^{\mu_1\mu_2\mu_3} = a^{\mu_1} g_\perp^{\mu_2\mu_3} + a^{\mu_2} g_\perp^{\mu_1\mu_3} + a^{\mu_3} g_\perp^{\mu_2\mu_1}; (a^2 b^2)^{\mu_1\mu_2\mu_3\mu_4} = a^{\mu_1} a^{\mu_2} b^{\mu_3} b^{\mu_4} + ... + b^{\mu_1} b^{\mu_2} a^{\mu_3} a^{\mu_4}; \\
(a^2 g_\perp)^{\mu_1\mu_2\mu_3\mu_4} = a^{\mu_1} a^{\mu_2} g_\perp^{\mu_2\mu_3} + a^{\mu_1} g_\perp^{\mu_1\mu_3} a^{\mu_4} + ... + g_\perp^{\mu_1\mu_2} a^{\mu_3} a^{\mu_4}; \\
(g_\perp g_\perp)^{\mu_1 \cdots \mu_4} = g_\perp^{\mu_1\mu_2} g_\perp^{\mu_3\mu_4} + g_\perp^{\mu_1\mu_3} g_\perp^{\mu_2\mu_4} + g_\perp^{\mu_1\mu_4} g_\perp^{\mu_2\mu_3}. \qquad \text{(C4)}
\end{aligned}
$$

The relevant integrals needed within operation of averaging over polar angle are:

$$
\begin{aligned}
\varphi_1 =< \frac{1-c^2}{d} >= \frac{1}{b^2}\left(1 - \frac{1-b^2}{2b} L\right); \quad \varphi_2 =< \frac{(1-c^2)^2}{d^2} = \frac{1}{2b^4}\left[3 - b^2 - \frac{1-b^2}{2b}(3+b^2) L\right]; \\
\varphi_3 =< \frac{1}{d} >= \frac{1}{b} L; \varphi_4 =< \frac{c^2(1-c^2)}{d^2} >= \frac{1}{b^4}[-\frac{3}{2} + \frac{1}{4}(7-3b^2)\frac{1}{b} L\Big]; \\
\varphi_5 =< \frac{1}{d^2} >= \frac{1}{1-b^2} + \frac{1}{2b} L; \quad \varphi_6 =< \frac{c^2}{d^2} >= \frac{1}{b^2}[\frac{1}{1-b^2} - \frac{1}{2b} L]; \quad L = \ln\frac{1+b}{1-b}. \qquad \text{(C5)}
\end{aligned}
$$

For the charged π-meson case we have

$$
\begin{aligned}
a_H^\pi &= 4 + 4\beta^2 \lambda \varphi_1 + \beta^4 \lambda^2 \frac{1}{2} \varphi_2; \\
a_G^\pi = a_F^\pi &= \frac{1}{2} \beta^4 \lambda^2 \varphi_2; \\
p^2 a_b^\pi = p^2 a_e^\pi &= -4 + \beta^2 \lambda(\varphi_1 - 2\varphi_2 \lambda) + \varphi_3 \lambda(-5 + \rho); \\
p^2 a_c^\pi = p^2 a_d^\pi &= \frac{1}{2} \beta^2 \lambda^2 (\rho - 1) b^2 \varphi_4; \\
p^2 a_a^\pi &= \frac{1}{2} \beta^2 \eta^2 \lambda^2 b^2 \varphi_4; \\
p^2 a_f^\pi &= \frac{1}{2} \beta^2 \sigma^2 \lambda^2 b^2 \varphi_4; \\
p^4 a_h^\pi &= 4 + 2\lambda(1-\rho)\varphi_3 + (16 + (1-\rho)^2)\lambda^2 \varphi_5. \qquad \text{(C6)}
\end{aligned}
$$

---

[1] R. Karplus and M. Neuman, Phus.REv. 80,(1950),380.
[2] V. Costantini, B. De-Tollis and G. Pistoni, Nuovo Cimento, 2A(1971),733.
[3] E. Kuraev and S. Sannikov, ZETP,44(1963),1015.
[4] A. I. Akhiezer and V. B. Berestetskij, Quantum Electrodynamics,GIZ, Moscow,1959.